

## Chapter 415

# Multivariate Analysis of Variance (MANOVA)

## Introduction

Multivariate analysis of variance (MANOVA) is an extension of common analysis of variance (ANOVA). In ANOVA, differences among various group means on a single-response variable are studied. In MANOVA, the number of response variables is increased to two or more. The hypothesis concerns a comparison of vectors of group means. When only two groups are being compared, the results are identical to Hotelling's  $T^2$  procedure.

The multivariate extension of the F-test is not completely direct. Instead, several test statistics are available, such as Wilks' Lambda and Lawley's trace. The actual distributions of these statistics are difficult to calculate, so we rely on approximations based on the F-distribution.

## Technical Details

A MANOVA has one or more factors (each with two or more levels) and two or more dependent variables. The calculations are extensions of the general linear model approach used for ANOVA.

Unlike the univariate situation in which there is only one statistical test available (the F-ratio), the multivariate situation provides several alternative statistical tests. We will describe these tests in terms of two matrices,  $H$  and  $E$ .  $H$  is called the *hypothesis matrix* and  $E$  is the *error matrix*. These matrices may be computed using a number of methods. In *NCSS*, we use the standard general linear models (GLM) approach in which a sum of squares and cross-products matrix is computed. This matrix is based on the dependent variables and independent variables generated for each degree of freedom in the model. It may be partitioned according to the terms in the model.

## MANOVA Test Statistics

For a particular p-variable multivariate test, assume that the matrices  $H$  and  $E$  have  $h$  and  $e$  degrees of freedom, respectively. Four tests may be defined as follows. See Seber (1984) for details. Let  $\theta_i$ ,  $\phi_i$ , and  $\lambda_i$  be the eigenvalues of  $H(E+H)^{-1}$ ,  $HE^{-1}$ , and  $E(E+H)^{-1}$  respectively. Note that these eigenvalues are related as follows:

$$\theta_i = 1 - \lambda_i = \frac{\phi_i}{1 + \phi_i}$$

$$\phi_i = \frac{\theta_i}{1 - \theta_i} = \frac{1 - \lambda_i}{\lambda_i}$$

$$\lambda_i = 1 - \theta_i = \frac{1}{1 + \phi_i}$$

## Multivariate Analysis of Variance (MANOVA)

### Wilks' Lambda

Define Wilks' Lambda as follows:

$$\Lambda_{p,h,e} = \frac{|E|}{|E+H|}$$

$$= \prod_{j=1}^p (1 - \theta_j)$$

with  $e \geq p$ .

The following approximation based on the F-distribution is used to determine significance levels:

$$F_{ph,ft-g} = \frac{(ft-g)(1-\Lambda^{1/t})}{ph\Lambda^{1/t}}$$

where

$$f = e - \frac{1}{2}(p-h+1)$$

$$g = \frac{ph-2}{2}$$

$$t = \begin{cases} \sqrt{\frac{p^2h^2-4}{p^2+h^2-5}} & \text{if } p^2+h^2-5 > 0 \\ 1 & \text{otherwise} \end{cases}$$

This approximation is exact if  $p$  or  $h \geq 2$ .

### Lawley - Hotelling Trace

The trace statistic,  $T_g^2$ , is defined as follows:

$$T_g^2 = e \sum_{j=1}^s \phi_j$$

where

$$s = \min(p, h)$$

The following approximation based on the F-distribution is used to determine significance levels:

$$F_{a,b} = \frac{T_g^2}{ce}$$

where

$$a = ph$$

$$b = 4 + (a+2)/(B-1)$$

$$c = \frac{a(b-2)}{b(e-p-1)}$$

## Multivariate Analysis of Variance (MANOVA)

$$B = \frac{(e + h - p - 1)(e - 1)}{(e - p - 3)(e - p)}$$

### Pillai's Trace

Pillai's trace statistic,  $V^{(s)}$ , is defined as follows:

$$V^{(s)} = \sum_{j=1}^s \theta_j = \text{tr}(H(E + H)^{-1})$$

where

$$s = \min(p, h)$$

The following approximation based on the F-distribution is used to determine significance levels:

$$F_{s(2m+s+1), s(2n+s+1)} = \frac{(2n + s + 1)V^{(s)}}{(2m + s + 1)(s - V^{(s)})}$$

where

$$s = \min(p, h)$$

$$m = (|p - h| - 1)/2$$

$$n = (e - p - 1)/2$$

### Roy's Largest Root

Roy's largest root,  $\phi_{\max}$ , is defined as the largest of the  $\phi_i$ 's. The following approximation based on the F-distribution is used to determine significance levels:

$$F_{(2\nu_1+2), (2\nu_2+2)} = \frac{2\nu_2 + 2}{2\nu_1 + 2} \phi_{\max}$$

where

$$s = \min(p, h)$$

$$\nu_1 = (|p - h| - 1)/2$$

$$\nu_2 = (e - p - 1)/2$$

---

## Which Test to Use

When the hypothesis degrees of freedom,  $h$ , is one, all four test statistics will lead to identical results. When  $h > 1$ , the four statistics will usually lead to the same result. When they do not, the following guidelines from Tabachnick (1989) may be of some help.

Wilks' Lambda, Lawley's trace, and Roy's largest root are often more powerful than Pillai's trace if  $h > 1$  and one dimension accounts for most of the separation among groups. Pillai's trace is more robust to departures from assumptions than the other three.

Tabachnick (1989) provides the following checklist for conducting a MANOVA. We suggest that you consider these issues and guidelines carefully.

---

## Assumptions and Limitations

The following assumptions are made when using a MANOVA.

1. The response variables are continuous.
2. The residuals follow the multivariate-normal probability distribution with means equal to zero.
3. The variance-covariance matrices of each group of residuals are equal.
4. The individuals are independent.

---

## Multivariate Normality and Outliers

MANOVA is robust to modest amount of skewness in the data. A sample size that produces 20 degrees of freedom in the univariate F-test is adequate to ensure robustness. Non-normality caused by the presence of outliers can cause severe problems that even the robustness of the test will not overcome. You should screen your data for outliers and run it through various univariate and multivariate normality tests and plots to determine if the normality assumption is reasonable.

---

## Homogeneity of Covariance Matrices

MANOVA makes the assumption that the within-cell (group) covariance matrices are equal. If the design is balanced so that there is an equal number of observations in each cell, the robustness of the MANOVA tests is guaranteed. If the design is unbalanced, you should test the equality of covariance matrices using Box's M test. If this test is significant at less than .001, there may be severe distortion in the alpha levels of the tests. You should only use Pillai's trace criterion in this situation.

---

## Linearity

MANOVA assumes linear relationships among the dependent variables within a particular cell. You should study scatter plots of each pair of dependent variables using a different color for each level of a factor. Look carefully for curvilinear patterns and for outliers. The occurrence of curvilinear relationships will reduce the power of the MANOVA tests.

---

## Multicollinearity and Singularity

*Multicollinearity* occurs when one dependent variable is almost a weighted average of the others. This collinearity may only show up when the data are considered one cell at a time. The R<sup>2</sup>-Other Y's in the Within-Cell Correlations Analysis report lets you determine if multicollinearity is a problem. If this R<sup>2</sup> value is greater than .99 for any variable, you should take corrective action (remove one of the variables). To correct for multicollinearity, begin removing the variables one at a time until all of the R<sup>2</sup>'s are less than .99. Do not remove them all at once! *Singularity* is the extreme form of multicollinearity in which the R<sup>2</sup> value is one.

Forms of multicollinearity may show up when you have very small cell sample sizes (when the number of observations is less than the number of variables). In this case, you must reduce the number of dependent variables.

---

## Data Structure

The data must be entered in a format that places the dependent variables and values of each factor side by side. An example of the data for a MANOVA design is shown in the table below. In this example, *WRATR* and *WRATA* are

## Multivariate Analysis of Variance (MANOVA)

the two dependent variables. *Treatment* and *Disability* are two factor variables. This database is stored in the file MANOVA1.

### MANOVA1 dataset (subset)

WRATR	WRATA	Treatment	Disability
115	108	1	1
98	105	1	1
107	98	1	1
90	92	2	1
85	95	2	1
80	81	2	1
100	105	1	2
105	95	1	2
95	98	1	2
70	80	2	2

---

## Unequal Sample Size and Missing Data

You should begin by screening your data. Pay particular attention to patterns of missing values. When using MANOVA, you should have more observations per factor category than you have dependent variables so that you can test the equality of covariance matrices using Box's M test.

NCSS ignores rows with missing values. If it appears that most of the missing values occur in one or two variables, you might want to leave these out of the analysis in order to obtain more data and hence more power.

NCSS uses the GLM procedure for calculating the hypothesis and error matrices. Each matrix is calculated as if it were fit last in the model. This is the recommended way of obtaining these matrices. This method is valid even when the sample sizes for the various groups are unequal.

---

## Procedure Options

This section describes the options available in this procedure.

---

### Variables Tab

These options control which variables are used in the analysis.

---

#### Response Variables

##### Response Variables

Specifies the response (dependent) variables to be analyzed.

---

#### Factor Specification

##### Factor Variable (1-10)

At least one factor variable must be specified. This variable's values indicates how the values of the response variable should be categorized. Examples of factor variables are gender, age groups, 'yes' or 'no' responses, etc. Note that the values in the variable may be either numeric or text. The treatment of text variables is specified for each variable by the Data Type option on the data base.

## Multivariate Analysis of Variance (MANOVA)

### Type

This option specifies whether the factor is fixed or random.

- **Fixed**

The factor includes all possible levels, like male and female for gender, includes representative values across the possible range of values, like low, medium, and high temperatures, or includes a set of values to which inferences will be limited, like New York, California, and Maryland.

- **Random**

The factor is one in which the chosen levels represent a random sample from the population of values. For example, you might select four classes from the hundreds in your state or you might select ten batches from an industrial process. The key is that a random sample is chosen. In *NCSS*, a random factor is “crossed” with other random and fixed factors. Two factors are crossed when each level of one includes all levels of the other.

---

### Model Specification

This section specifies the experimental design model.

#### Which Model Terms

A design in which main effect and interaction terms are included is called a saturated model. Often, it is useful to omit various interaction terms from the model. This option lets you specify which interactions to keep very easily. If the selection provided here is not flexible enough for your needs, you can specify custom here and enter the model directly.

The options included are as follows.

- **Full Model**

The complete, saturated model is analyzed. This option requires that you have no missing cells, although you can have an unbalanced design. Hence, you cannot use this option with Latin square or fractional factorial designs.

- **Up to 1-Way**

A main-effects only model is run. All interactions are omitted.

- **Up to 2-Way**

All main-effects and two-way interactions are included in the model.

- **Up to 3-Way**

All main-effects, two-way, and three-way interactions are included in the model.

- **Up to 4-Way**

All main-effects, two-way, three-way, and four-way interactions are included in the model.

- **Custom**

This option indicates that you want the Custom Model (given in the next box) to be used.

#### Custom Model

When a Custom Model is selected (see Which Model Terms), the model itself is entered here. If all main effects and interactions are desired, you can enter the word “ALL” here. For complicated designs, it is usually easier to check the next option, Write Model in ‘Custom Model’ Field, and run the procedure. The appropriate model will be generated and placed in this box. You can then edit it as you desire.

## Multivariate Analysis of Variance (MANOVA)

The model is entered using letters separated by the plus sign. For example, a three-factor factorial in which only two-way interactions are needed would be entered as follows:

A+B+AB+C+AC+BC

Note that repeated-measures designs are not allowed.

### Write Model in 'Custom Model' Field

When this option is checked, no analysis is performed when the procedure is run. Instead, a copy of the full model is stored in the Custom Model box. You can then delete selected terms from the model without having to enter all the terms you want to keep.

---

## Reports Tab

The following options control which reports are displayed.

---

### Select Reports

#### EMS Report ... Univariate F's

Specify whether to display the indicated reports.

#### Test Alpha

The value of alpha for the statistical tests and power analysis. Usually, this number will range from 0.1 to 0.001. A common choice for alpha is 0.05, but this value is a legacy from the age before computers when only printed tables were available. You should determine a value appropriate for your particular study.

---

### Report Options

#### Precision

Specify the precision of numbers in the report. Single precision will display seven-place accuracy, while double precision will display thirteen-place accuracy.

#### Variable Names

Indicate whether to display the variable names or the variable labels.

#### Value Labels

Indicate whether to display the data values or their labels.

---

## Plots Tab

The following options specify the plot(s) of group means.

---

### Select Plots

#### Means Plot(s) and Subject Plot

Specify whether to display the indicated plots. Click the plot format button to change the plot settings.

#### Y-Axis Scaling

Specify the method for calculating the minimum and maximum along the vertical axis. *Separate* means that each plot is scaled independently. *Uniform* means that all plots use the overall minimum and maximum of the data. This option is ignored if a minimum or maximum is specified.

## Example 1 – Multivariate Analysis of Variance

This section presents an example of how to run an analysis of the data contained in the MANOVA1 dataset. You may follow along here by making the appropriate entries or load the completed template **Example 1** by clicking on Open Example Template from the File menu of the Multivariate Analysis of Variance (MANOVA) window.

### 1 Open the MANOVA1 dataset.

- From the File menu of the NCSS Data window, select **Open Example Data**.
- Click on the file **MANOVA1.NCSS**.
- Click **Open**.

### 2 Open the MANOVA window.

- Using the Analysis menu or the Procedure Navigator, find and select the **Multivariate Analysis of Variance (MANOVA)** procedure.
- On the menus, select **File**, then **New Template**. This will fill the procedure with the default template.

### 3 Specify the variables.

- On the Multivariate Analysis of Variance (MANOVA) window, select the **Variables tab**.
- Double-click in the **Response Variable** box. This will bring up the variable selection window.
- Select **WRATR** and **WRATA** from the list of variables and then click **Ok**. “WRATR-WRATA” will appear in the Response Variable box.
- Double-click in the **first Factor Variable** box. This will bring up the variable selection window.
- Select **Treatment** from the list of variables and then click **Ok**. “Treatment” will appear in the first Factor Variable box.
- Double-click in the **second Factor Variable** box. This will bring up the variable selection window.
- Select **Disability** from the list of variables and then click **Ok**. “Disability” will appear in the second Factor Variable box.

### 4 Run the procedure.

- From the Run menu, select **Run Procedure**. Alternatively, just click the green Run button.

## Expected Mean Squares Section

### Expected Mean Squares Section

Source Term	DF	Term Fixed?	Denominator Term	Expected Mean Square
A (Treatment)	1	Yes	S(AB)	S+bsA
B (Disability)	2	Yes	S(AB)	S+asB
AB	2	Yes	S(AB)	S+sAB
S(AB)	12	No		S

Note: Expected Mean Squares are for the balanced cell-frequency case.

The Expected Mean Square expressions are provided to show the appropriate error term for each factor. The correct error term for a factor is that term that is identical except for the factor being tested.

### Source Term

The source of variation or term in the model.

### DF

The degrees of freedom. The number of observations “used” by this term.



## Multivariate Analysis of Variance (MANOVA)

### Term Fixed?

Indicates whether the term is “fixed” or “random.”

### Denominator Term

Indicates the term used as the denominator in the F-ratio.

### Expected Mean Square

This expression represents the expected value of the corresponding mean square if the design was completely balanced. “S” represents the expected value of the mean square error (sigma). The uppercase letters represent either the adjusted sum of squared treatment means if the factor is fixed, or the variance component if the factor is random. The lowercase letter represents the number of levels for that factor, and “s” represents the number of replications of the experimental layout.

These EMS expressions are provided to determine the appropriate error term for each factor. The correct error term for a factor is that term whose EMS is identical except for the factor being tested.

## MANOVA Tests Section

Manova Tests Section						
Term(DF) Test Statistic	Test Value	DF1	DF2	F-Ratio	Prob Level	Decision (0.05)
<b>A(1):Treatment</b>						
Wilks' Lambda	0.137721	2	11	34.44	0.000018	Reject
Hotelling-Lawley Trace	6.261036	2	11	34.44	0.000018	Reject
Pillai's Trace	0.862279	2	11	34.44	0.000018	Reject
Roy's Largest Root	6.261036	2	11	34.44	0.000018	Reject
WRATR	2090.88889	1	12	46.12	0.000019	Reject
WRATA	1494.22222	1	12	33.25	0.000089	Reject
<b>B(2):Disability</b>						
Wilks' Lambda	0.255263	4	22	5.39	0.003528	Reject
Hotelling-Lawley Trace	2.895034	4	20	7.24	0.000896	Reject
Pillai's Trace	0.750481	4	24	3.60	0.019460	Reject
Roy's Largest Root	2.887241	2	12	17.32	0.000290	Reject
WRATR	260.388889	2	12	5.74	0.017784	Reject
WRATA	563.388889	2	12	12.54	0.001151	Reject
<b>AB(2)</b>						
Wilks' Lambda	0.908068	4	22	0.27	0.893037	Accept
Hotelling-Lawley Trace	0.100954	4	20	0.25	0.904790	Accept
Pillai's Trace	0.092192	4	24	0.29	0.881598	Accept
Roy's Largest Root	0.098039	2	12	0.59	0.570550	Accept
WRATR	1.055556	2	12	0.02	0.977029	Accept
WRATA	26.388889	2	12	0.59	0.571116	Accept

This report gives the results of the various significance tests. Usually, the four multivariate tests will lead to the same conclusions. When they do not, refer to the discussion of these tests found earlier in this chapter. Once a multivariate test has found a term significant, use the univariate ANOVA to determine which of the variables and factors are “causing” the significance.

### Term(DF)

The term in the design model with the degrees of freedom of the term in parentheses.

### Test Statistic

The name of the statistical test shown on this row of the report. The four multivariate tests are followed by the univariate F-tests of each variable.

### Test Value

The value of the test statistic.

## Multivariate Analysis of Variance (MANOVA)

### DF1

The numerator degrees of freedom of the F-ratio corresponding to this test.

### DF2

The denominator degrees of freedom of the F-ratio corresponding to this test.

### F-Ratio

The value of the F-test corresponding to this test. In some cases, this is an exact test. In other cases, this is an approximation to the exact test. See the discussion of each test to determine if it is exact or approximate.

### Prob Level

The significance level of the above F-ratio. The probability of an F-ratio larger than that obtained by this analysis. For example, to test at an alpha of 0.05, this probability would have to be less than 0.05 to make the F-ratio significant.

### Decision(0.05)

The decision to accept or reject the null hypothesis at the given level of significance. Note that you specify the level of significance when you select Alpha.

---

## Within Correlations\Covariances Section

### Within Correlations\Covariances Section

	WRATR	WRATA
WRATR	45.33333	2.583333
WRATA	0.0572313	44.94444

This report displays the correlations and covariances formed by averaging across all the individual group covariance matrices. The correlations are shown in the lower-left half of the matrix. The within-group covariances are shown on the diagonal and in the upper-right half of the matrix.

---

## Within-Cell Correlations Analysis Section

### Within-Cell Correlations Analysis

Variable	R-Squared Other Y's	Canonical Variate	Eigenvalue	Percent of Total	Cumulative Total
Wratr	0.003275	1	1.057231	52.86	52.86
Wrata	0.003275	2	0.942769	47.14	100.00

This report analyzes the within-cell correlation matrix. It lets you diagnose multicollinearity problems as well as determine the number of dimensions that are being used. This is useful in determining if Pillai's trace should be used.

### R-Squared Other Y's

This is the R-Squared index of this variable with the other variables. When this value is larger than 0.99, severe multicollinearity problems exist. If this happens, you should remove the variable with the largest R-Squared and re-run your analysis.

### Canonical Variate

The identification numbers of the canonical variates that are generated during the analysis. The total number of variates is the smaller of the number of variables and the number of degrees of freedom in the model.

## Multivariate Analysis of Variance (MANOVA)

### Eigenvalue

The eigenvalues of the within correlation matrix. Note that this value is not associated with the variable at the beginning of the row, but rather with the canonical variate number directly to the left.

### Percent of Total

The percent that the eigenvalue is of the total. Note that the sum of the eigenvalues will equal the number of variates. If the percentage accounted for by the first eigenvalue is relatively large (70 or 80 percent), Pillai's trace will be less powerful than the other three multivariate tests.

### Cumulative Total

The cumulative total of the Percent of Total column.

---

## Univariate Analysis of Variance Section

### Analysis of Variance Table for WRATR

Source Term	DF	Sum of Squares	Mean Square	F-Ratio	Prob Level	Power (Alpha=0.05)
A (Treatment)	1	2090.889	2090.889	46.12	0.000019*	0.999988
B (Disability)	2	520.7778	260.3889	5.74	0.017784*	0.763859
AB	2	2.111111	1.055556	.02	0.977029	0.052757
S	12	544	45.33333			
Total (Adjusted)	17	3157.778				
Total	18					

\* Term significant at alpha = 0.05

This is the standard ANOVA report as documented in the General Linear Models chapter. A separate report is displayed for each of the dependent variables.

---







## Means and Plots Section

### Means and Standard Errors of WRATR

Term	Count	Mean	Standard Error
All	18	89.11111	
A: Treatment			
1	9	99.88889	2.234687
2	9	78.33334	2.234687
B: Disability			
1	6	95.83334	2.736922
2	6	88.83334	2.736922
3	6	82.66666	2.736922
AB: Treatment,Disability			
1,1	3	106.6667	3.870592
1,2	3	100	3.870592
1,3	3	93	3.870592
2,1	3	85	3.870592
2,2	3	77.66666	3.870592
2,3	3	72.33334	3.870592

## Multivariate Analysis of Variance (MANOVA)

### Plots Section



This report provides the least-squares means and standard errors for each variable. Note that the standard errors are calculated from the mean square error of the ANOVA table. They are not the standard errors that would be calculated from the individual cells.