

Chapter 142

Non-Inferiority Tests for the Ratio of Two Within-Subject Variances in a $2 \times 2M$ Replicated Cross- Over Design

Introduction

This procedure calculates power and sample size of *non-inferiority* tests of within-subject variabilities from a $2 \times 2M$ replicated cross-over design. This routine deals with the case in which the statistical hypotheses are expressed in terms of the ratio of the within-subject variances.

This design is used to compare two treatments which are administered to subjects in different orders. It has two treatment sequences. Here, M is the number of times a particular treatment is received by a subject. For example, if $M = 2$, the design is a 2×4 cross-over. The two sequences would often be

sequence 1: R T R T

sequence 2: T R T R

It is assumed that either there is no carry-over from one measurement to the next, or there is an ample washout period between measurements.

Technical Details

This procedure uses the formulation given in Chow, Shao, Wang, and Lokhnygina (2018).

Suppose x_{ijkl} is the response in the i th sequence ($i = 1, 2$), j th subject ($j = 1, \dots, N_i$), k th treatment ($k = T, C$), and l th replicate ($l = 1, \dots, M$). The mixed effect model analyzed in this procedure is

$$x_{ijkl} = \mu_k + \gamma_{ikl} + S_{ijk} + e_{ijkl}$$

where μ_k is the k th treatment effect, γ_{ikl} is the fixed effect of the l th replicate on treatment k in the i th sequence, S_{ij1} and S_{ij2} are random effects of the ij th subject, and e_{ijkl} is the within-subject error term which is normally distributed with mean 0 and variance $V_k = \sigma_{Wk}^2$

Unbiased estimators of these variances are found after applying an orthogonal transformation matrix P to the x 's as follows

$$z_{ijk} = P' x_{ijk}$$

where P is an $m \times m$ matrix such that $P'P$ is diagonal and $\text{var}(z_{ijkl}) = \sigma_{Wk}^2$.

For example, in a 2x4 cross-over design the z 's become

$$z_{ijk1} = \frac{x_{ijk1} + x_{ijk2}}{2} = \bar{x}_{ijk}$$

and

$$z_{ijk2} = \frac{x_{ijk1} - x_{ijk2}}{\sqrt{2}} = \bar{x}_{ijk}$$

In this case, the within-subject variances are estimated as

$$\hat{V}_T = \frac{1}{(N_1 + N_2 - 2)(M - 1)} \sum_{i=1}^2 \sum_{j=1}^{N_i} \sum_{l=1}^M (z_{ijTl} - \bar{z}_{i.Tl})^2$$

and

$$\hat{V}_C = \frac{1}{(N_1 + N_2 - 2)(M - 1)} \sum_{i=1}^2 \sum_{j=1}^{N_i} \sum_{l=1}^M (z_{ijCl} - \bar{z}_{i.Cl})^2$$

Testing Non-Inferiority

The following hypotheses are usually used to test for non-inferiority

$$H_0: \frac{\sigma_{WT}^2}{\sigma_{WC}^2} \geq R_0 \text{ versus } H_A: \frac{\sigma_{WT}^2}{\sigma_{WC}^2} < R_0,$$

where R_0 is the non-inferiority limit.

The corresponding test statistic is $T = (\hat{V}_1/\hat{V}_2)/R_0$.

Upon making the usual normality assumptions, T is distributed as an $F_{d,d}$ random variable where

$$d = (N_1 + N_2 - 2)(M - 1).$$

Power

The power of this combination of tests is given by

$$\text{Power} = P\left(F < \frac{R0}{R1} F_{\alpha,d,d}\right)$$

where F is the common F distribution with the indicated degrees of freedom, α is the significance level, and $R1$ is the value of the variance ratio stated by the alternative hypothesis. Lower quantiles of F are used in the equation.

A simple binary search algorithm can be applied to this power function to obtain an estimate of the necessary sample size.

Procedure Options

This section describes the options that are specific to this procedure. These are located on the Design tab. For more information about the options of other tabs, go to the Procedure Window chapter.

Design Tab

The Design tab contains the parameters associated with this test such as sample sizes, alpha, and power.

Solve For

Solve For

This option specifies the parameter to be solved for from the other parameters. Under most situations, you will select either *Power* or *Sample Size*.

Power and Alpha

Power

This option specifies one or more values for power. Power is the probability of rejecting a false null hypothesis, and is equal to one minus Beta. Beta is the probability of a type-II error, which occurs when a false null hypothesis is not rejected. In this procedure, a type-II error occurs when you fail to reject the null hypothesis of nonequivalent means when in fact the means are equivalent.

Values must be between zero and one. Historically, the value of 0.80 (Beta = 0.20) was used for power. Now, 0.90 (Beta = 0.10) is also commonly used.

A single value may be entered here or a range of values such as *0.8 to 0.95 by 0.05* may be entered.

Alpha

This option specifies one or more values for the probability of a type-I error. A type-I error occurs when a true null hypothesis is rejected. In this procedure, a type-I error occurs when you reject the null hypothesis of non-equivalence when in fact the null hypothesis is true.

Values must be between zero and one. Historically, the value of 0.05 has been used for alpha. This means that about one test in twenty will falsely reject the null hypothesis. You should pick a value for alpha that represents the risk of a type-I error you are willing to take in your experimental situation.

You may enter a range of values such as *0.01 0.05 0.10* or *0.01 to 0.10 by 0.01*.

Sample Size (When Solving for Sample Size)

Sequence Allocation

Select the option that describes the constraints on $N1$ or $N2$ or both.

The options are

- **Equal ($N1 = N2$)**

This selection is used when you wish to have equal sample sizes in each sequence. Since you are solving for both sample sizes at once, no additional sample size parameters need to be entered.

- **Enter $N2$, solve for $N1$**

Select this option when you wish to fix $N2$ at some value (or values), and then solve only for $N1$. Please note that for some values of $N2$, there may not be a value of $N1$ that is large enough to obtain the desired power.

- **Enter $R = N2/N1$, solve for $N1$ and $N2$**

For this choice, you set a value for the ratio of $N2$ to $N1$, and then PASS determines the needed $N1$ and $N2$, with this ratio, to obtain the desired power. An equivalent representation of the ratio, R , is

$$N2 = R * N1.$$

- **Enter percentage in Sequence 1, solve for $N1$ and $N2$**

For this choice, you set a value for the percentage of the total sample size that is in Sequence 1, and then PASS determines the needed $N1$ and $N2$ with this percentage to obtain the desired power.

$N2$ (Sample Size, Sequence 2)

This option is displayed if Sequence Allocation = "Enter $N2$, solve for $N1$ "

$N2$ is the number of items or individuals sampled from the Sequence 2 population.

$N2$ must be ≥ 2 . You can enter a single value or a series of values.

R (Sequence Sample Size Ratio)

This option is displayed only if Sequence Allocation = "Enter $R = N2/N1$, solve for $N1$ and $N2$."

R is the ratio of $N2$ to $N1$. That is,

$$R = N2 / N1.$$

Use this value to fix the ratio of $N2$ to $N1$ while solving for $N1$ and $N2$. Only sample size combinations with this ratio are considered.

$N2$ is related to $N1$ by the formula:

$$N2 = [R \times N1],$$

where the value $[Y]$ is the next integer $\geq Y$.

For example, setting $R = 2.0$ results in a Sequence 2 sample size that is double the sample size in Sequence 1 (e.g., $N1 = 10$ and $N2 = 20$, or $N1 = 50$ and $N2 = 100$).

R must be greater than 0. If $R < 1$, then $N2$ will be less than $N1$; if $R > 1$, then $N2$ will be greater than $N1$. You can enter a single or a series of values.

Non-Inferiority Tests for the Ratio of Two Within-Subject Variances in a 2x2M Rep. Cross-Over Design**Percent in Sequence 1**

This option is displayed only if Sequence Allocation = "Enter percentage in Sequence 1, solve for N1 and N2."

Use this value to fix the percentage of the total sample size allocated to Sequence 1 while solving for $N1$ and $N2$. Only sample size combinations with this Sequence 1 percentage are considered. Small variations from the specified percentage may occur due to the discrete nature of sample sizes.

The Percent in Sequence 1 must be greater than 0 and less than 100. You can enter a single or a series of values.

Sample Size (When Not Solving for Sample Size)**Sequence Allocation**

Select the option that describes how individuals in the study will be allocated to Sequence 1 and to Sequence 2.

The options are

- **Equal ($N1 = N2$)**
This selection is used when you wish to have equal sample sizes in each sequence. A single per sequence sample size will be entered.
- **Enter N1 and N2 individually**
This choice permits you to enter different values for $N1$ and $N2$.
- **Enter N1 and R, where $N2 = R * N1$**
Choose this option to specify a value (or values) for $N1$, and obtain $N2$ as a ratio (multiple) of $N1$.
- **Enter total sample size and percentage in Sequence 1**
Choose this option to specify a value (or values) for the total sample size (N), obtain $N1$ as a percentage of N , and then $N2$ as $N - N1$.

Sample Size Per Sequence

This option is displayed only if Sequence Allocation = "Equal ($N1 = N2$)."

The Sample Size Per Sequence is the number of items or individuals sampled. Since the sample sizes are the same in each sequence, this value is the value for $N1$, and also the value for $N2$.

The Sample Size Per Sequence must be ≥ 2 . You can enter a single value or a series of values.

N1 (Sample Size, Sequence 1)

*This option is displayed if Sequence Allocation = "Enter N1 and N2 individually" or "Enter N1 and R, where $N2 = R * N1$."*

$N1$ is the number of items or individuals sampled from the Sequence 1 population.

$N1$ must be ≥ 2 . You can enter a single value or a series of values.

N2 (Sample Size, Sequence 2)

This option is displayed only if Sequence Allocation = "Enter N1 and N2 individually."

$N2$ is the number of items or individuals sampled from the Sequence 2 population.

$N2$ must be ≥ 2 . You can enter a single value or a series of values.

Non-Inferiority Tests for the Ratio of Two Within-Subject Variances in a 2x2M Rep. Cross-Over Design**R (Sequence Sample Size Ratio)**

*This option is displayed only if Sequence Allocation = "Enter N1 and R, where $N2 = R * N1$."*

R is the ratio of $N2$ to $N1$. That is,

$$R = N2/N1$$

Use this value to obtain $N2$ as a multiple (or proportion) of $N1$.

$N2$ is calculated from $N1$ using the formula:

$$N2 = [R \times N1],$$

where the value $[Y]$ is the next integer $\geq Y$.

For example, setting $R = 2.0$ results in a Sequence 2 sample size that is double the sample size in Sequence 1.

R must be greater than 0. If $R < 1$, then $N2$ will be less than $N1$; if $R > 1$, then $N2$ will be greater than $N1$. You can enter a single value or a series of values.

Total Sample Size (N)

This option is displayed only if Sequence Allocation = "Enter total sample size and percentage in Sequence 1."

This is the total sample size, or the sum of the two sequence sample sizes. This value, along with the percentage of the total sample size in Sequence 1, implicitly defines $N1$ and $N2$.

The total sample size must be greater than one, but practically, must be greater than 3, since each sequence sample size needs to be at least 4.

You can enter a single value or a series of values.

Percent in Sequence 1

This option is displayed only if Sequence Allocation = "Enter total sample size and percentage in Sequence 1."

This value fixes the percentage of the total sample size allocated to Sequence 1. Small variations from the specified percentage may occur due to the discrete nature of sample sizes.

The Percent in Sequence 1 must be greater than 0 and less than 100. You can enter a single value or a series of values.

M (Number of Replicates)

Enter one or more values for M: the number of replicates in a $2 \times 2M$ replicated cross-over design. M is the number of times each treatment is measured on each subject.

For example, $M = 2$ refers to a 2×4 replicated cross-over design. The two sequences used for this design are often

Sequence 1: R T R T

Sequence 2: T R T R.

You can enter a single value such as 2, a series of values such as 2 3 4, or 2 to 8 by 1.

The range is $M \geq 2$.

Effect Size

R0 (Non-Inferiority Variance Ratio)

Enter one or more values for the non-inferiority limit for the ratio of the two within-subject variances. When the ratio of the sample variances is less than this value, the treatment group is concluded to be “non-inferior” to the control group.

This value must be greater than one. Popular choices are 1.5 and 2.

R1 (Actual Variance Ratio)

Enter one or more values for within-subject variance ratio assumed by the alternative hypothesis. This is the value of $\sigma^2_{WT} / \sigma^2_{WC}$ at which the power is calculated.

This value should be less than R0.

Example 1 – Finding Sample Size

A company has developed a generic drug for treating rheumatism and wants to show that it is non-inferior to the standard drug in terms of the within-subject variability. A 2 x 4 replicated cross-over design will be used to test the non-inferiority.

Company researchers set the non-inferiority limit to 1.5, the significance level to 0.05, the power to 0.90, M to 2, and the actual variance ratio values between 0.8 and 1.2. They want to investigate the range of required sample size values assuming that the two group sample sizes are equal.

Setup

This section presents the values of each of the parameters needed to run this example. First, from the PASS Home window, load the **Non-Inferiority Tests for the Ratio of Two Within-Subj Variances in a 2x2M Rep. Cross-Over Design** procedure window. You may then make the appropriate entries as listed below, or open **Example 1** by going to the **File** menu and choosing **Open Example Template**.

<u>Option</u>	<u>Value</u>
Design Tab	
Solve For	Sample Size
Power	0.90
Alpha	0.05
Sequence Allocation	Equal (N1 = N2)
M (Number of Replicates)	2
R0 (Non-Inferiority Variance Ratio)	1.5
R1 (Actual Variance Ratio)	0.8 0.9 1 1.1 1.2

Annotated Output

Click the Calculate button to perform the calculations and generate the following output.

Numeric Results

Actual Power	Sequence 1 Sample Size N1	Sequence 2 Sample Size N2	N	Number of Replicates M	Non- Inferiority Variance Ratio R0	Actual Variance Ratio R1	Alpha
0.9013	45	45	90	2	1.500	0.800	0.050
0.9036	68	68	136	2	1.500	0.900	0.050
0.9009	106	106	212	2	1.500	1.000	0.050
0.9007	180	180	360	2	1.500	1.100	0.050
0.9004	346	346	692	2	1.500	1.200	0.050

References

- Chow, S.C., Shao, J., Wang, H., and Lohknygina, Y. 2018. Sample Size Calculations in Clinical Research, Third Edition. Taylor & Francis/CRC. Boca Raton, Florida.
- Chow, S.C. and Liu, J.P. 2014. Design and Analysis of Clinical Trials, Third Edition. John Wiley & Sons. Hoboken, New Jersey.

Non-Inferiority Tests for the Ratio of Two Within-Subject Variances in a 2x2M Rep. Cross-Over Design

Report Definitions

Actual Power is the actual power achieved. Because N_1 and N_2 are discrete, this value is usually slightly larger than the target power.

N_1 is the number of subjects in sequence 1.

N_2 is the number of subjects in sequence 2.

N is the total number of subjects. $N = N_1 + N_2$.

M is the number of replicates. That is, it is the number of times a treatment measurement is repeated on a subject.

R_0 is the non-inferiority limit for the within-subject variance ratio.

R_1 is the value of the within-subject variance ratio at which the power is calculated.


Alpha is the probability of rejecting a true null hypothesis, H_0 .

Summary Statements

A study is being conducted to assess the non-inferiority of the within-subject variability of two groups by looking at the ratio of their variances. If the sample variance ratio falls below the non-inferiority limit, the treatment (group 1) is concluded to be 'non-inferior' to the reference group. Otherwise, it is not. Sequence sample sizes of 45 and 45 achieve 90% power to reject the null hypothesis of inferiority at a significance level of 0.050. The non-inferiority limit is 1.500. The variance ratio at which the power is calculated is 0.800. The number of times each treatment measurement is repeated on a subject is 2.

This report gives the sample sizes for the indicated scenarios.

Plot Section



These plots show the relationship between sample size and R_1 .

Example 2 – Validation using Chow et al. (2018)

The following example of finding the sample size for a 2x4 cross-over design is shown in Chow *et al.* (2018) pages 199 - 200.

Find the sample size when the non-inferiority limit is 1.21, the significance level to 0.05, M is 2, the power is 0.8, and the alternative variance ratio value 0.4444. They obtained $N_1 = N_2 = 14$.

Setup

This section presents the values of each of the parameters needed to run this example. First, from the PASS Home window, load the **Non-Inferiority Tests for the Ratio of Two Within-Subj Variances in a 2x2M Rep. Cross-Over Design** procedure window. You may then make the appropriate entries as listed below, or open **Example 2** by going to the **File** menu and choosing **Open Example Template**.

<u>Option</u>	<u>Value</u>
Design Tab	
Solve For	Sample Size
Power.....	0.80
Alpha.....	0.05
Sequence Allocation	Equal (N1 = N2)
M (Number of Replicates).....	2
R0 (Non-Inferiority Variance Ratio)	1.21
R1 (Actual Variance Ratio)	0.444

Output

Click the Calculate button to perform the calculations and generate the following output.

Numeric Results								
Actual Power	Sequence 1 Sample Size N1	Sequence 2 Sample Size N2	N	Number of Replicates M	Non-Inferiority Variance Ratio R0	Actual Variance Ratio R1	Alpha	
0.8079	14	14	28	2	1.210	0.444	0.050	

The sequence sample sizes are 14 and match the results of Chow et al. (2018).